

RECOMMENDED LITERATURE FOR FOREIGN STUDENTS

1. Ang M, Fenwick E, Wong TY, Lamoureux E, Luo N. Utility of EQ-5D to assess patients undergoing cataract surgery. *Optom Vis Sci* 2013;90:861–6.
2. Bailey IL, Lovie-Kitchin JE. Visual acuity testing: from the laboratory to the clinic. *Vision Res* 2013;90:2–9.
3. Brown MM, Brown GC, Sharma S, Busbee B, Brown H. Quality of life associated with unilateral and bilateral good vision. *Ophthalmology* 2001;108:643–7.
4. Clark A, Ng JQ, Morlet N, Tropiano E, Mahendran P, Spilsbury K, et al. Quality of life after postoperative endophthalmitis. *Clin Experiment Ophthalmol* 2008;36:526–31.
5. Colquitt JL, Jones J, Tan SC, Takeda AL, Clegg AJ, Price A. Ranibizumab and pegaptanib for the treatment of age-related macular degeneration: a systematic review and economic evaluation. *Health Technol Assess* 2008;12(16).
6. De Juan-Marcos L, Blanco-Blanco JF, Hernandez-Galilea E. Visual function and quality of life in pseudophakic patients before and after capsulotomy. *Eur J Ophthalmol* 2012;22:943–9.
7. Dolders MG, Nijkamp MD, Nuijts RM, van den Borne B, Hendrikse F, Ament A, et al. Cost effectiveness of foldable multifocal intraocular lenses compared to foldable monofocal intraocular lenses for cataract surgery. *Br J Ophthalmol* 2004;88:1163–8.
8. Edmunds B, Thompson JR, Salmon JF, Wormald RP. The National Survey of Trabeculectomy. III. Early and late complications. *Eye*. 2002;16(3):297–303.
9. Harper R, Radi N, Reeves BC, Fenerty C, Spencer AF, Batterbury M. Agreement between ophthalmologists and optometrists in optic disc assessment: training implications for glaucoma co-management. *Graefes Arch Clin Exp Ophthalmol*. 2001;239(5):342–50.
10. Jefferis JM, Taylor J-P, Collerton J, Jagger C, Kingston A, Davies K, et al. The association between diagnosed glaucoma and cataract and cognitive

performance in very old people: cross-sectional findings from the Newcastle 85+ study. *Ophthalmic Epidemiol* 2013;20:82–8.

- 11.Kaushik S, Pandav SS, Ram J. Neuroprotection in glaucoma. *J Postgrad Med.* 2003;49(1):90–5.
- 12.Kishimoto F, Naito T, Hasebe S, Ohtsuki H. Time trade-off utility analysis for surgical intervention in comitant strabismus, glaucoma, and cataract. *Acta Med Okayama* 2012;66:191–201.
- 13.Naeim A, Keeler EB, Gutierrez PR, Wilson MR, Reuben D, Mangione CM. Is cataract surgery cost-effective among older patients with a low predicted probability for improvement in reported visual functioning? *Med Care* 2006;44:982–9.
- 14.Neudorfer M, Sadetzki S, Anisimova S, Geyer O. Nonpenetrating deep sclerectomy with the use of adjunctive mitomycin C. *Ophthalmic surgery, Lasers & Imaging.* 2004;35(1):6–12.
- 15.Rosser DA, Laidlaw DA, Murdoch IE. The development of a ‘reduced logMAR’ visual acuity chart for use in routine clinical practice. *Br J Ophthalmol* 2001;85:432–6.
- 16.Schneck ME, Hagerström-Portnoy G, Lott LA, Brabyn JA. Ocular contributions to age-related loss in coarse stereopsis. *Optom Vis Sci* 2013;77:531–6.
- 17.Szymanski A, Gierek-Lapinska A, Koziak M, Gierek-Ciaciura S. A fluorophotometric study of corneal endothelium after trabeculectomy using different concentrations of Mitomycin-C. *International Ophthalmology.* 1997;20(1–3):95–9.
- 18.Theodossiades J, Murdoch I. What optic disc parameters are most accurately assessed using the direct ophthalmoscope. *Eye.* 2001;15(Pt 3):283–7.
- 19.Thomas R, George T, Braganza A, Muliyl J. The flashlight test and van Herick's test are poor predictors for occludable angles. *Australian and New Zealand Journal of Ophthalmology.* 1996;24(3):251–6.

20. Tosh J, Brazier J, Evans P, Longworth L. A review of generic preference-based measures of health-related quality of life in visual disorders. *Value Health* 2012;15:118–27.
21. Tsai J-C, Chang HW. Comparison of the effects of brimonidine 0.2% and timolol 0.5% on retinal nerve fibre layer thickness in ocular hypertensive patients: A prospective, unmasked study. *Journal of Ocular Pharmacology & Therapeutics*. 2005;21(6):475–82.
22. Vass C, Findl O, Sycha T, Bauer P, Schmetterer L. Medical interventions for primary open angle glaucoma and ocular hypertension. *Cochrane Database Syst Rev*. 2004;(3):CD003167.
23. Vetrugno M, Cardascia N, Cantatore F, Sborgia C. Comparison of the effects of bimatoprost and timolol on intraocular pressure and pulsatile ocular blood flow in patients with primary open-angle glaucoma: A prospective, open-label, randomized, two-arm, parallel-group study. *Current Therapeutic Research, Clinical & Experimental*. 2004;65(6):444–54.
24. Weiss KB, Sullivan SD. The health economics of asthma and rhinitis. I. Assessing the economic impact. *J Allergy Clin Immunol*. 2001;107(1):3–8.
25. Wilkins M, Indar A, Wormald R. Intra-operative mitomycin C for glaucoma surgery. *Cochrane Database Syst Rev*. 2005;(4):CD002897.
26. Yalvac IS, Sahin M, Eksioglu U, Midillioglu IK, Aslan BS, Duman S. Primary viscocanalostomy versus trabeculectomy for primary open-angle glaucoma: three-year prospective randomized clinical trial.[see comment] *Journal of Cataract & Refractive Surgery*. 2004;30(10):2050–7.
27. Zhou Z, Althin R, Sforzolini BS, Dhawan R. Persistency and treatment failure in newly diagnosed open angle glaucoma patients in the United Kingdom. *British Journal of Ophthalmology*. 2004;88(11):1391–4.